

MH907: Diabetes in Pregnancy

[View Online](#)


Bell, R., Glinianaia, S. V., Tennant, P. W. G., Bilous, R. W., & Rankin, J. (2012). Peri-conception hyperglycaemia and nephropathy are associated with risk of congenital anomaly in women with pre-existing diabetes: a population-based cohort study. *Diabetologia*, 55(4), 936–947. <https://doi.org/10.1007/s00125-012-2455-y>

Bellamy, L., Casas, J.-P., Hingorani, A. D., & Williams, D. (2009). Type 2 diabetes mellitus after gestational diabetes: a systematic review and meta-analysis. *The Lancet*, 373(9677), 1773–1779. [https://doi.org/10.1016/S0140-6736\(09\)60731-5](https://doi.org/10.1016/S0140-6736(09)60731-5)

Confidential Enquiry into Maternal and Child Health. (n.d.). Pregnancy in women with type 1 and type 2 diabetes 2002–2003: England, Wales and Northern Ireland. CEMACH: Confidential Enquiry into Maternal and Child Health. <http://www.hqip.org.uk/national-programmes/a-z-of-clinical-outcome-review-programmes/cemach-reports/>

Global Library of Women's Medicine. (n.d.). <http://www.glowm.com/>

Great Britain. (2003). National service framework for diabetes: delivery strategy [Electronic resource]. Department of Health.

Kim, S. Y., England, J. L., Sharma, J. A., & Njoroge, T. (2011). Gestational Diabetes Mellitus and Risk of Childhood Overweight and Obesity in Offspring: A Systematic Review. *Experimental Diabetes Research*, 2011, 1–9. https://warwick.summon.serialssolutions.com/2.0.0/link/0/eLvHCXMwjV1JS8NAFB5sD-JF3K1LGfAcm2WjyBShFGsRS7FV8RYms2ApptKk-A_83b5Z7KFehBxnJvCS977vzdsQisJr39uwCXGheCRYIhZAHwQlsqBck2ESRzGVpm54ME0ejtnjW_bk8mWrvxF9AL-ORqxOrG1v1kAN0BHtog_ffs2v7gdm6t8S0B7AJuKK8ja2At19107v12yDTxpcGeyhXUclcc9-wX20JcsDtD1yle9D9N39Ymv57b2sandvh10WS4VHuptmvaowKwWezKo5Xii8blaMxybpWXvfZoGbAYBnJR4rZQOy3Y45_Qbbt_TwdN3ZGduwgVtxhF4Gd8_9oeGJ3icEL_20oQRlBlgFSwkwCoo9_XwPclJvQEJCWKB0oBiGVJJ CglPhg6rlIVSh1JY9ExapaLUp4izHkiAPUpUwUHb5pmshChH3NQfTiV-S105WSZf9oWGblxLei41xLPrcRb6MTKeb0ljGUCBDU4-9f-c7Rj73P1c4Ga9XIL3WI1VLPnG-jRv910Da_wg9OJbDP

Learn about diabetes using interactive tools. (n.d.). <https://web.archive.org/web/20140618224054/http://diabetes.niddk.nih.gov/resources/HealthTools/index.aspx>

Meltzer, S. J. (2010). Prepregnancy Care: A Shared Responsibility. *Diabetes Care*, 33(12), 2713–2715. <https://0-care-diabetesjournals-org.pugwash.lib.warwick.ac.uk/content/33/12/2713>

Miller, E., Hare, J. W., Cloherty, J. P., Dunn, P. J., Gleason, R. E., Soeldner, J. S., & Kitzmiller,

- J. L. (1981). Elevated Maternal Hemoglobin A1C in Early Pregnancy and Major Congenital Anomalies in Infants of Diabetic Mothers. *New England Journal of Medicine*, 304(22), 1331–1334.
<http://0-search.proquest.com.pugwash.lib.warwick.ac.uk/docview/1870344756?accountid=14888>
- Murphy, H. R., Roland, J. M., Skinner, T. C., Simmons, D., Gurnell, E., Morrish, N. J., Soo, S.-C., Kelly, S., Lim, B., Randall, J., Thompsett, S., & Temple, R. C. (2010). Effectiveness of a Regional Prepregnancy Care Program in Women With Type 1 and Type 2 Diabetes: Benefits beyond glycemic control. *Diabetes Care*, 33(12), 2514–2520.
<https://0-care-diabetesjournals-org.pugwash.lib.warwick.ac.uk/content/33/12/2514>
- Murphy, H. R., Temple, R. C., Ball, V. E., Roland, J. M., Steel, S., Zill-E-Huma, R., Simmons, D., Royce, L. R., & Skinner, T. C. (2010). Personal experiences of women with diabetes who do not attend pre-pregnancy care. *Diabetic Medicine*, 27(1), 92–100.
<https://doi.org/10.1111/j.1464-5491.2009.02890.x>
- National Diabetes Support Team. (n.d.).
<http://webarchive.nationalarchives.gov.uk/20130316063827/http://www.diabetes.nhs.uk/>
National Institute for Clinical Excellence. (2008). Insulin Pump Therapy: Continuous subcutaneous insulin infusion for the treatment of diabetes mellitus. National Institute for Clinical Excellence. <https://www.nice.org.uk/guidance/ta151>
- National Institute for Clinical Excellence. (2015). Diabetes in pregnancy: management of diabetes and its complications from conception to the postnatal period: NICE guideline. National Institute for Clinical Excellence. <https://www.nice.org.uk/guidance/ng3>
- National Service Framework. (n.d.).
<http://webarchive.nationalarchives.gov.uk/+www.nhs.uk/nhsengland/NSF/pages/Nationalserviceframeworks.aspx>
- NHS Evidence. (n.d.). <https://www.evidence.nhs.uk/>
- NICE - National Institute for Health and Care Excellence. (n.d.). <http://www.nice.org.uk/>
- Pearson, D., Kernaghan, D., Lee, R., & Penney, G. (2006). Short communication: The relationship between pre-pregnancy care and early pregnancy loss, major congenital anomaly or perinatal death in type I diabetes mellitus. *BJOG: An International Journal of Obstetrics & Gynaecology*, 114(1), 104–107.
<https://doi.org/10.1111/j.1471-0528.2006.01145.x>
- Poor glycated haemoglobin control and adverse pregnancy outcomes in type 1 and type 2 diabetes mellitus: Systematic review of observational studies. (2006). *BMC Pregnancy and Childbirth*, 6. <https://0-pubmed-ncbi-nlm-nih-gov.pugwash.lib.warwick.ac.uk/17074087/>
- Ray, J. G., O'Brien, T. E., & Chan, W. S. (2001). Preconception care and the risk of congenital anomalies in the offspring of women with diabetes mellitus: a meta-analysis. *QJM*, 94(8), 435–444. <https://doi.org/10.1093/qjmed/94.8.435>
- Saravanan, P., & Yajnik, C. S. (2010). Role of maternal vitamin B12 on the metabolic health of the offspring: a contributor to the diabetes epidemic? *The British Journal of Diabetes & Vascular Disease*, 10(3), 109–114. <https://doi.org/10.1177/1474651409358015>

Scottish Intercollegiate Guidelines Network. (2010). Management of diabetes: a national clinical guideline: Vol. SIGN publication [Electronic resource]. SIGN.
<https://www.sign.ac.uk/assets/sign116.pdf>